

**ANRS - AC 11 : RESISTANCE GROUP
GENOTYPE INTERPRETATION: NUCLEOSIDE AND NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS**

	Mutations associated to resistance	Mutations associated to « possible resistance »
ZDV	<ul style="list-style-type: none"> • T215Y/F • At least 3 mutations among : M41L, D67N, K70R, L210W, T215A/C/D/E/G/H/I/L/N/S/V, K219Q/E [1, 2, 3, 4] • Q151M • Insertion at codon 69 	<ul style="list-style-type: none"> • T215A/C/D/E/G/H/I/L/N/S/V [1, 2, 3, 4]
3TC/FTC	<ul style="list-style-type: none"> • M184V/I • Insertion at codon 69 	<ul style="list-style-type: none"> • K65R [11, 12, 16] • Q151M
ddl	<ul style="list-style-type: none"> • At least a score of + 2 among: M41L + T69D + L74V/I + 215Y/F + K219Q/E – K70R – M184 V/I [5, 14, 15, 17, 18] • L74V/I without any mutations among M41L, T69D, K70R, M184 V/I, T215Y/F, K219Q/E [19] • Q151M • Insertion at codon 69 	<ul style="list-style-type: none"> • K65R [11, 12]
d4T	<ul style="list-style-type: none"> • V75A/M/S/T • T215Y/F [6] • At least 3 mutations among : M41L, D67N, K70R, L210W, T215A/C/D/E/G/H/I/L/N/S/V, K219Q/E [4, 7, 14, 15] • Q151M • Insertion at codon 69 	<ul style="list-style-type: none"> • T215A/C/D/E/G/H/I/L/N/S/V [4, 7]
ABC	<ul style="list-style-type: none"> • At least 5 mutations among : M41L, D67N, L74V/I, M184V/I, L210W, T215Y/F [8, 19, 29] • No TAMs and at least 2 mutations among: K65R, L74V/I, Y115F and M184V/I [24, 25, 26, 27, 28, 29] • Q151M • Insertion at codon 69 	<ul style="list-style-type: none"> • 4 mutations among : M41L, D67N, L74V/I, M184V/I, L210W, T215Y/F [8, 19, 29] • K65R [9, 11, 12]
TDF	<ul style="list-style-type: none"> • At least 6 mutations among: M41L, E44D, D67N, T69D/N/S, L74V, L210W, T215Y/F [13, 20] • K65R [9, 10, 11, 12] • Insertion at codon 69 	<ul style="list-style-type: none"> • 3, 4 or 5 mutations among: M41L, E44D, D67N, T69D/N/S, L74V, L210W, T215Y/F [13] • K70E [21, 22, 23]

TAMs = M41L, D67N, K70R, L210W, T215Y/F, K219Q/E

ZDV: zidovudine, 3TC: lamivudine, FTC: emtricitabine, ddl: didanosine, d4T: stavudine, ABC: abacavir, TDF: tenofovir

Mutations associated to antiretroviral regimen but with uncertain signification: E44A, D67E/G, T69A, K219N/R

**ANRS - AC 11 : RESISTANCE GROUP
GENOTYPE INTERPRETATION: NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS**

	Mutations associated to resistance	Mutations associated to « possible resistance »
EFV	<ul style="list-style-type: none"> • L100I • K101E • K103H/N/S/T [1] • V106M [2] • Y181C/I • Y188C/L • G190A/C/E/Q/S/T/V • P225H • M230L 	
NVP	<ul style="list-style-type: none"> • A98S (for HIV-1 subtype C only) [3] • L100I • K101E • K103H/N/S/T [1] • V106A/M [2] • Y181C/I • Y188C/H/L • G190A/C/E/Q/S/T/V • M230L 	<ul style="list-style-type: none"> • A98S (for other subtypes than HIV-1 subtype C) [3]
ETV	<ul style="list-style-type: none"> • At least 4 among: V90I, A98G, L100I, K101E/P, V106I, V179D/F, Y181C/I/V, G190A/S [4] 	<ul style="list-style-type: none"> • 3 mutations among: V90I, A98G, L100I, K101E/P, V106I, V179D/F, Y181C/I/V, G190A/S [4]

EFV: efavirenz, NVP: nevirapine, ETV: etravirine

Mutations associated to antiretroviral regimen but with uncertain signification: K101H/N/Q

ANRS - AC 11 : RESISTANCE GROUP

GENOTYPE INTERPRETATION: PROTEASE INHIBITORS

	Mutations associated to resistance	Mutations associated to « possible resistance »
IDV	<ul style="list-style-type: none"> • M46I/L • V82A/F/M/S/T [11] • I84A/V [8] • L90M and at least 2 among : K20M/R, L24I, V32I, M36I, I54V/L/M/T, A71V/T, G73S/A, V77I 	<ul style="list-style-type: none"> • L90M
SQV/RTV 1000/100 mg BID	<ul style="list-style-type: none"> • G48V • At least 4 mutations among: L10F/I/M/R/V, I15A/V, K20I/M/R/T, L24I, I62V, G73S/T, V82A/F/S/T, I84V, L90M [9] 	<ul style="list-style-type: none"> • 3 mutations among: L10F/I/M/R/V, I15A/V, K20I/M/R/T, L24I, I62V, G73S/T, V82A/F/S/T, I84V, L90M [9]
NFV	<ul style="list-style-type: none"> • D30N • I84A/V [8] • N88S/D • L90M 	<ul style="list-style-type: none"> • V82A/F/S/T and at least 2 among: L10I, M36I, M46I/L, I54V/L/M/T, A71V/T, V77I [1]
FPV/RTV 700/100 mg BID	<ul style="list-style-type: none"> • I50V • V32I and I47A/V [2, 13, 14] • At least 4 mutations among: L10F/I/V, L33F, M36I, I54A/L/M/S/T/V, I62V, V82A/C/F/G, I84V, L90M [2, 20] 	
LPV/r	<ul style="list-style-type: none"> • At least 6 mutations among : L10F/I/R/V, K20M/R, L24I, L33F, M46I/L, I50V, F53L, I54M/L/T/V, L63P, A71I/L/V/T, V82A/F/S/T, I84V, L90M [3, 4, 5, 21] • I47A [15, 16] • L76V [18, 19] 	<ul style="list-style-type: none"> • 4 or 5 mutations among: L10F/I/R/V, K20M/R, L24I, L33F, M46I/L, I50V, F53L, I54M/L/T/V, L63P, A71I/L/V/T, V82A/F/S/T, I84V, L90M [3, 4, 5, 21]
ATV/RTV 300/100 mg QD	<ul style="list-style-type: none"> • I50L [6] • At least 3 mutations among: L10F/I/V, G16E, L33F/I/V, M46I/L, D60E, I84V, I85V, L90M [7, 12, 22] 	
TPV/RTV 500/200 mg BID	<ul style="list-style-type: none"> • At least a score of + 3* : 36I/L/V – 53L/W/Y + 58E + 69I/K/N/Q/R/Y + 89I/M/R/T/V [10, 23] 	<ul style="list-style-type: none"> • A score of + 2* : 36I/L/V – 53L/W/Y + 58E + 69I/K/N/Q/R/Y + 89I/M/R/T/V [10, 23]
DRV/RTV 600/100 mg BID	<ul style="list-style-type: none"> • At least 4 mutations among: V11I, V32I, L33F, I47V, I50V, I54L/M, T74P, L76V, I84V, L89V [17, 24, 25, 26] 	<ul style="list-style-type: none"> • 3 mutations among: V11I, V32I, L33F, I47V, I50V, I54L/M, T74P, L76V, I84V, L89V [17, 24, 25, 26]

July 2008- Version n°17

**IDV: indinavir, SQV: saquinavir, NFV: nelfinavir, RTV: ritonavir, FPV: fosamprenavir, LPV: lopinavir, ATV:atazanavir, TPV: tipranavir,
DRV : darunavir**

Mutations associated to antiretroviral regimen but with uncertain signification: K20T, M46V

N88S mutation is associated to better virological response to fosamprenavir [2]

* Insufficient data for HIV-1 subtype non-B

ANRS - AC 11 : RESISTANCE GROUP
GENOTYPE INTERPRETATION: FUSION INHIBITOR

Mutations associated to resistance	
ENF T20	<ul style="list-style-type: none">• G36A/D/E/S/V [1, 2, 3, 4, 5, 6, 7]• V38A/E/K/M• Q40H/K/P/T• N42D/T• N43D/H/K/S• L44M• L45Q/M

ENF (T20): enfuvirtide

ANRS - AC 11 : RESISTANCE GROUP
GENOTYPE INTERPRETATION: INTEGRASE INHIBITORS

	Mutations associated to resistance
RAL	<ul style="list-style-type: none">• Q148H/G/K/R/E [1, 2]• N155H [1, 2]• E157Q [2]• E92Q [1, 2]• Y143C/H/R [3, 4, 5]

RAL: raltegravir

July 2008- Version n°17

**ANRS - AC 11 : RESISTANCE GROUP
GENOTYPE INTERPRETATION FOR HIV-2**

NUCLEOSIDE AND NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS
<ul style="list-style-type: none">• Q151M : Resistance to d4T, ABC [1]• Q151M + M184V : Resistance to all NRTI except TDF [2]• M184V : Resistance to 3TC/FTC [2]
NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS
<ul style="list-style-type: none">• Naturally resistant to all NNRTI [3, 4]
PROTEASE INHIBITORS
<ul style="list-style-type: none">• Naturally resistant to APV and fosAPV [3, 4]• Contradictory data for ATV , TPV [3, 5]
FUSION INHIBITOR
<ul style="list-style-type: none">• Naturally resistant to T20 [3, 4]
INTEGRASE INHIBITORS
<ul style="list-style-type: none">• Q148K/R : Resistance to raltegravir [6]• N155H : Resistance to raltegravir [7]

REFERENCES

Nucleoside and nucleotide Reverse Transcriptase Inhibitors

- 1/ Riva C et al. Transmitted virus with substitutions at position 215 and risk of virological failure in antiretroviral naive patients starting highly active antiretroviral therapy. XI International HIV drug resistance workshop : basic principles and clinical implications , 2-5 July 2002, Seville, Spain, abstract 124.
- 2/ Lanier ER et al. Effect of baseline 215D/C/S « revertant » mutations on virological response to lamivudine /zidovudine- containing regimens and emergence of 215Y upon virological failure. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain abstract 146.
- 3/ Garcia-Lerma et al. Increased ability for selection of zidovudine resistance in a distinct class of wild-type HIV-1 from drug-naive persons. PNAS 2001 ; 98 : 13907-13912.
- 4/ Chappey C et al. Evolution of amino acid 215 in HIV-1 reverse transcriptase in response to intermittent drug selection. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 32.
- 5/ Costagliola D et al. Presence of thymidine –associated mutations and response to d4T, abacavir and ddi in the control arm of the Narval ANRS 088 trial. 8th Conference on Retroviruses and opportunistic Infections. 4-8 february 2001, Chicago, Illinois, poster 450.
- 6/ Calvez V et al. Impact of stavudine phenotype and thymidine analogs mutations on viral response to stavudine plus lamivudine in ALTIS 2 ANRS trial. Antiviral Therapy 2002, 7(3):211-218.

July 2008- Version n°17

- 7/ García-Lerma JG, MacInnes H, Bennett D, Reid P, Nidtha S, Weinstock H, Kaplan JE, Heneine W. A novel genetic pathway of human immunodeficiency virus type 1 resistance to stavudine mediated by the K65R mutation. *J Virol.* 2003 May;77(10):5685-93.
- 8/ Brun-Vézinet F et al. Clinically relevant interpretation of genotype for resistance to abacavir : a study from the Narval trial (ANRS 088). *AIDS* 2003; 17(12):1795-802.
- 9/ Stone C, Ait-Khaled M, Craig C, Griffin P, Tisdale M. Human immunodeficiency virus type 1 reverse transcriptase mutation selection during in vitro exposure to tenofovir alone or combined with abacavir or lamivudine. *Antimicrob Agents Chemother.* 2004 Apr;48(4):1413-5.
- 10/ Miller MD et al. Multivariate analyses of antiviral response to tenofovir DF therapy in antiretroviral-experienced patients. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain abstract 14.
- 11/ Margot NA, Lu B, Cheng A, Miller MD; Study 903 Team. Resistance development over 144 weeks in treatment-naive patients receiving tenofovir disoproxil fumarate or stavudine with lamivudine and efavirenz in Study 903. *HIV Med.* 2006 Oct;7(7):442-50.
- 12/ Parikh et al. K65R : a multi-nucleoside resistance mutation of a low but increasing frequency. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 136.
- 13/ Masquelier B et al. Genotypic and pharmacological determinants of the virological response to tenofovir in nucleoside reverse transcriptase inhibitor-experienced patients. *Antivir Ther.* 2004 ; 9(3):315-23.
- 14/ Izopet J. et al. Mutations conferring resistance to zidovudine diminish the antiviral effect of stavudine plus didanosine. *Journal of Medical Virology* 1999 ; 59 : 507-511.

July 2008- Version n°17

- 15/ Pellegrin I. et al. Emergence of zidovudine and multidrug-resistance mutations in the HIV-1 reverse transcriptase gene in therapy-naive patients receiving stavudine plus didanosine combination therapy. STADI group. AIDS 1999 ; 13 : 1705-1709.
- 16/ Mulamba GB et al. Pre-steady state kinetic analysis of the incorporation of FTC 5'-monophosphate and 3TC 5'-monophosphate by mutants HIV-1 RTs K65R, K65R/Q151M and Q151M. 16th International Conference on Antiviral Research, 27 April-1 May 2003, Savannah, USA. Abstract 39.
- 17/ Marcelin AG et al. Clinically relevant genotype interpretation of resistance to didanosine. Antimicrob Agents Chemother. 2005 May;49(5):1739-44.
- 18/ Capdepon S et al. An additive/subtractive genotypic score as a determinant of the virological response to didanosine in HIV-1 infected patients. J Clin Virol. 2006 May;36(1): 36-42.
- 19/ Assoumou L, Brun-Vézinet F, Cozzi-Lepri A, Kuritzkes D, Phillips A, Zolopa A, Degruittola V, Miller V, Costagliola D; Standardization and Clinical Relevance of HIV Drug Resistance Testing Project of the Forum for Collaborative HIV Research. Initiatives for Developing and Comparing Genotype Interpretation Systems: External Validation of Existing Systems for Didanosine against Virological Response. J Infect Dis. 2008 Jul 3.
- 20/ Waters J et al. K65R, L74V and TAMs in HIV-1 RT associated with reduced response to tenofovir DF in antiretroviral-experienced patients. 13th Conference on Retroviruses and opportunistic Infections. 5-8 february 2006, Denver, Colorado, poster 633.
- 21/ Kagan RM, Lee TS, Ross L, Lloyd RM Jr, Lewinski MA, Potts SJ. Molecular basis of antagonism between K70E and K65R tenofovir-associated mutations in HIV-1 reverse transcriptase. Antiviral Res. 2007 Sep;75(3):210-8.

July 2008- Version n°17

22/ Kagan R et al. Adefovir HIV-1 RT mutation K70E in the age of tenofovir. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 93.

23/ Sluis-Cremer N, Sheen CW, Zelina S, Torres PS, Parikh UM, Mellors JW. Molecular mechanism by which the K70E mutation in human immunodeficiency virus type 1 reverse transcriptase confers resistance to nucleoside reverse transcriptase inhibitors. *Antimicrob Agents Chemother.* 2007 Jan;51(1):48-53.

24/ Miller V et al. HIV-1 reverse transcriptase (RT) genotype and susceptibility to RT inhibitors during abacavir monotherapy and combination therapy. *AIDS* 2000; 14:163–171.

25/ Moyle GJ, et al. Ziagen Once-Daily in Antiretroviral Combination Therapy (CNA30021) Study Team. Abacavir once or twice daily combined with once-daily lamivudine and efavirenz for the treatment of antiretroviral-naïve HIV-infected adults: results of the Ziagen Once Daily in Antiretroviral Combination Study. *J Acquir Immune Defic Syndr.* 2005 Apr 1;38(4):417-25.

26/ Irlbeck D *et al.* Treatment-emergent mutations for previously naïve HIV-infected adults failing ZDV+3TC+EFV and ABC+3TC+EFV (CNA30024). *11th Conference on Retroviruses and Opportunistic Infections.* San Francisco, CA, USA. February 8–11, 2004. Abstract 661.

27/. Rodriguez-French A, *et al.* The NEAT study: a 48-week open-label study to compare the antiviral efficacy and safety of GW433908 versus nelfinavir in antiretroviral therapy-naïve HIV-1-infected patients. *J Acquir Immune Defic Syndr* 2004; **35**:22–32.

28/. Gathe JC Jr et al. SOLO: 48-week efficacy and safety comparison of once-daily fosamprenavir /ritonavir versus twice-daily nelfinavir in naïve HIV-1-infected patients. *AIDS.* 2004 Jul 23;18(11):1529-37

July 2008- Version n°17

29/ Wirden et al. Risk factors for selection of the L74I reverse transcriptase mutation in human immunodeficiency virus type 1-infected patients. *Antimicrob Agents Chemother.* 2006 Jul;50(7):2553-6.

Non nucleoside transcriptase inhibitors

1/ Harrigan PR, Mo T, Wynhoven B, Hirsch J, Brumme Z, McKenna P, Pattery T, Vingerhoets J, Bachelier LT. Rare mutations at codon 103 of HIV-1 reverse transcriptase can confer resistance to non-nucleoside reverse transcriptase inhibitors. *AIDS.* 2005 Mar 24;19(6):549-54.

2/ Brenner B et al. A V106M mutation in HIV-1 clade C viruses exposed to efavirenz confers cross-resistance to non-nucleoside reverse transcriptase inhibitors. *AIDS.* 2003 Jan 3;17(1):F1-5.

3/ Deshpande A et al. Resistance mutations in subtype C HIV type 1 isolates from Indian patients of Mumbai receiving NRTIs plus NNRTIs and experiencing a treatment failure: resistance to AR. *AIDS Res Hum Retroviruses* 2007; 23 : 335-40.

4/ Vingerhoets J et al. Impact of baseline NNRTI mutations on the virological response to TMC125 in the Phase III clinical trials DUET-1 and DUET-2. XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 32.

Protease inhibitors

1/ Phenotype/genotype correlation from Stanford database.

July 2008- Version n°17

- 2/ Masquelier B, Assoumou KL, Descamps D, Bocket L, Cottalorda J, Ruffault A, Marcelin AG, Morand-Joubert L, Tamalet C, Charpentier C, Peytavin G, Antoun Z, Brun-Vézinet F, Costagliola D; ANRS Resistance Study Group. Clinically validated mutation scores for HIV-1 resistance to fosamprenavir/ritonavir. *J Antimicrob Chemother.* 2008 Jun;61(6):1362-8.
- 3/ Marcelin AG, Cohen-Codar I, King MS, Colson P, Guillevic E, Descamps D, Lamotte C, Schneider V, Ritter J, Segondy M, Peigue-Lafeuille H, Morand-Joubert L, Schmuck A, Ruffault A, Palmer P, Chaix ML, Mackiewicz V, Brodard V, Izopet J, Cottalorda J, Kohli E, Chauvin JP, Kempf DJ, Peytavin G, Calvez V. Virological and pharmacological parameters predicting the response to lopinavir-ritonavir in heavily protease inhibitor-experienced patients. *Antimicrob Agents Chemother.* 2005 May;49(5):1720-6.
- 4/ Maillard A, Chapplain JM, Tribut O, Bentué-Ferrer D, Tattevin P, Arvieux C, Michelet C, Ruffault A. The use of drug resistance algorithms and genotypic inhibitory quotient in prediction of lopinavir-ritonavir treatment response in human immunodeficiency virus type 1 protease inhibitor-experienced patients. *J Clin Virol.* 2007 Feb;38(2):131-8.
- 5/ Masquelier B et al. Human Immunodeficiency virus type 1 genotypic and pharmacokinetic determinants of the virological response to lopinavir-ritonavir-containing therapy in protease inhibitor-experienced patients. *Antimicrob Agents and Chemother* 2002 ; 46 : 2926-2932.
- 6/ Colonna RJ et al. Identification of I50L as the signature atazanavir (ATV)-resistance mutation in treatment naïve HIV-1 infected patients receiving ATV-containing regimens. *JID* 2004; 189: 1802-10.
- 7/ Vora S et al. Clinical validation of atazanavir/ritonavir genotypic resistance score in PI-experienced patients. *AIDS* 2006 Jan 2;20(1):35-40.

July 2008- Version n°17

8/ Mo H, Parkin N, Stewart KD, Lu L, Dekhtyar T, Kempf DJ, Molla A. Identification and structural characterization of I84C and I84A mutations that are associated with high-level resistance to human immunodeficiency virus protease inhibitors and impair viral replication. *Antimicrob Agents Chemother.* 2007 Feb;51(2):732-5.

9/ Marcelin AG, Flandre P, de Mendoza C, Roquebert B, Peytavin G, Valer L, Wirden M, Abbas S, Katlama C, Soriano V, Calvez V. Clinical validation of saquinavir/ritonavir genotypic resistance score in protease-inhibitor-experienced patients. *Antivir Ther.* 2007;12(2):247-52.

10/ Marcelin AG et al. Tipranavir-ritonavir genotypic resistance score in protease inhibitor-experienced patients. *Antimicrob Agents Chemother.* 2008 Sept;52(9).

11/ Camacho R et al. Different substitutions under selective pressure at protease codon 82 in HIV-1 subtype G compared to subtype B infected individuals including a novel I82M resistance mutation. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 138.

12/ Marcelin AG et al. External validation of atazanavir/ritonavir genotypic score in HIV-1 protease inhibitor experienced patients. *JAIDS* 2006; 42 (1): 127-8.

13/ Brutus A et al. The Neat Study: Virologic response in subjects receiving GW433908 (908) BID by viral load/CD4 cell Counts at study entry. XV International AIDS Conference, July 11-16 2004, Bangkok ,Thailand, abstract TuPeB 4500.

July 2008- Version n°17

14/ Pellegrin I, Breilh D, Coureau G, Boucher S, Neau D, Merel P, Lacoste D, Fleury H, Saux MC, Pellegrin JL, Lazaro E, Dabis F, Thiébaud R; ANRS Co3 Aquitaine Cohort. Interpretation of genotype and pharmacokinetics for resistance to fosamprenavir-ritonavir-based regimens in antiretroviral-experienced patients. *Antimicrob Agents Chemother.* 2007 Apr;51(4):1473-80.

15/ Friend J et al. Isolated lopinavir resistance after virological rebound of a ritonavir/lopinavir-based regimen. *AIDS.* 2004 Sep 24;18(14):1965-6.

16/ de Mendoza C et al. Prevalence of the HIV-1 protease mutation I47A in clinical practice and association with lopinavir resistance. *AIDS* 2006 Apr 24; 20(7): 1071-4.

17/ de Meyer S, Vangeneugden T, van Baelen B, de Paepe E, van Marck H, Picchio G, Lefebvre E, de Béthune MP. Resistance profile of darunavir: combined 24-week results from the POWER trials. *AIDS Res Hum Retroviruses.* 2008 Mar;24(3):379-88.

18/ Nijhuis N et al. A novel genetic pathway involving L76V and M46I leading to lopinavir/r resistance. XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 127.

19/ Delaugerre C et al. Protease gene mutations in a trial comparing first-line lopinavir/ritonavir monotherapy to lopinavir/ritonavir + zidovudine/lamivudine (MONARK Trial). XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 75.

20/ Marcelin AG et al. Genotypic analysis of the virological response to fosamprenavir/ritonavir in clinical trials: CONTEXT and TRIAD. 14th Conference on Retroviruses and opportunistic Infections. 25-28 february 2007, Los Angeles, California, poster 608.

July 2008- Version n°17

21/ Hill A et al. Re-evaluation of resistance algorithms for lopinavir/ritonavir in the TITAN trial. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 126.

22/ Di Giambenedetto S et al. A rigorous statistical learning method for the estimation and validation of weighted drug susceptibility scores applied to in vivo virological outcome prediction in atazanavir/ritonavir-containing HAART. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 95.

23/ Bethell R et al. No effect of subtype on susceptibility and virological response to TPV/r for treatment experienced patients. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 111.

24/ Descamps D et al. Mutations associated with virologic response to darunavir/ritonavir in HIV-1 infected PI-experienced patients. 11th European AIDS Conference EACS; 24-27 october 2007, Madrid, Spain, abstract PS4/3.

25/ De Meyer S. et al. Effect of baseline and on-treatment mutations on the antiretroviral activity of darunavir/ritonavir and lopinavir/ritonavir: results of a randomised, controlled, Phase III study (TITAN). 4th International AIDS Society Conference on HIV Pathogenesis, Treatment and Prevention; Sydney, Australia; 22–25 July 2007. Abstract WEPEB038.

26/ De Meyer S. et al. Phenotypic and genotypic determinants of resistance to darunavir: analysis of data from treatment-experienced patients in POWER 1, 2, 3 and DUET-1 and 2. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 31.

July 2008- Version n°17

Fusion inhibitor

1/ Sista P et al. Subgroup analysis of baseline susceptibility and early virological response to enfuvirtide in the combined TORO studies. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 55.

2/ Mink M et al. Impact of HIV-1 gp41 amino acid substitutions (position 36-45) on susceptibility to T20 (enfuvirtide) in vitro ; analysis of primary virus isolates recovered from patients during chronic enfuvirtide treatment and site-directed mutants in NL4-3. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain, abstract 22.

3/ Greenberg ML et al. Enfuvirtide (T-20) and T-1249 resistance : observations from phase II clinical trials of enfuvirtide in combination with oral antiretrovirals and a phase I/II dose-ranging monotherapy trial of T-1249. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain, abstract 128.

4/ Greenberg ML et al. Baseline and on-treatment susceptibility to enfuvirtide seen in TORO 1 and TORO 2 to 24 weeks. 10th Conference on Retroviruses and Opportunistic Infections, February 10-14, 2003, Boston, USA, abstract 141.

5/ Chakraborty B et al. Replicative fitness of HIV-1 strains with reduced susceptibility to protease-, reverse transcriptase- and entry (enfuvirtide)-inhibitors. XIII International HIV drug resistance workshop : basic principles and clinical implications, 8-12 June 2004, Tenerife, Spain, abstract 61.

6/ Walmsley SL et al. Development of a new genotypic resistance assay involving the entire gp41 sequence for evaluating resistance to enfuvirtide. XIII International HIV drug resistance workshop : basic principles and clinical implications, 8-12 June 2004, Tenerife, Spain, abstract 134.

July 2008- Version n°17

7/ Melby T et al. Evolution of enfurvitide resistance in longitudinal samples obtained after continued enfuvirtide dosing post-virological failure. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 67.

Integrase inhibitors

1/ Hazuda DJ et al. Resistance to the HIV-integrase inhibitor raltegravir: analysis of protocol 005, a Phase II study in patients with triple-class resistant HIV-1 infection XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 8.

2/ Malet I et al. Mutations associated with failure of raltegravir treatment affect integrase sensitivity to the inhibitor in vitro. *Antimicrob Agents Chemother.* 2008 Apr;52(4):1351-8.

3/ Hatano H et al. Virological and immunological outcomes in a cohort of patients failing integrase inhibitors. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 10.

4/ Da Silva D et al. Mutational patterns in the HIV-1 integrase related to virological failures on raltegravir-containing regimens. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 12.

5/ Ceccherini-Silberstein F et al. Virological response and resistance in multi-experienced patients treated by raltegravir. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 18.

July 2008- Version n°17

HIV-2

- 1/ Damond F et al. In vitro phenotypic susceptibility to nucleoside reverse transcriptase inhibitors of HIV-2 isolates with the Q151M mutation in the reverse transcriptase. *Antivir Ther* 2005; 10(7): 861-5.
- 2/ Damond F et al. Phenotypic susceptibility to nucleoside reverse transcriptase inhibitors of HIV-2 isolates with the Q151M mutation in the reverse transcriptase. 13th Conference on Retroviruses and opportunistic Infections. 5-8 february 2006, Denver, Colorado, poster 607.
- 3/ Desbois D, Roquebert B, Peytavin G et al. In vitro phenotypic susceptibility of human immunodeficiency virus type 2 clinical isolates to protease inhibitors. *Antimicrob Agents Chemother*. 2008 Apr;52(4):1545-8.
- 4/ Witvrouw E et al. Susceptibility of HIV-2, SIV and SHIV to various anti-HIV-1 compounds: implications for treatment and postexposure prophylaxis. *Antivir Ther* 2004; 9(1): 57-65.
- 5/ Parkin NT et al. Antiretroviral drug resistance in non-subtype B, HIV-2 and SIV. *Antivir Ther* 2004; 9(1): 3-12.
- 6/Roquebert B, Blum L, Collin G et al. Selection of the Q148R integrase inhibitor resistance mutation in failing raltegravir containing regimen. *AIDS* 2008 *in press*.
- 7/ Garrett N, Xub L, Smit E, et al. Raltegravir treatment response in an HIV-2 infected patient: a case report. *AIDS* 2008, 22: 1091-1098.