

**ANRS - AC 11: RESISTANCE GROUP  
GENOTYPE INTERPRETATION: NUCLEOSIDE AND NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS**

	Mutations associated with resistance	Mutations associated with « possible resistance »
ZDV	<ul style="list-style-type: none"> <li>• T215A/C/D/E/G/H/I/L/N/S/V/Y/F [1, 2, 3, 4]</li> <li>• At least 3 mutations among: M41L, D67N, K70R, L210W, K219Q/E [1, 2, 3, 4]</li> <li>• Q151M</li> <li>• Insertion at codon 69</li> </ul>	
3TC/FTC	<ul style="list-style-type: none"> <li>• K65R [11, 12, 16]</li> <li>• M184V/I</li> <li>• Insertion at codon 69</li> </ul>	<ul style="list-style-type: none"> <li>• Q151M</li> </ul>
ddl	<ul style="list-style-type: none"> <li>• At least a score of + 2 among: M41L + T69D + 215Y/F + K219Q/E – K70R – M184 V/I [5, 14, 15, 17, 18]</li> <li>• K65R [11, 12]</li> <li>• L74V/I [19]</li> <li>• Q151M</li> <li>• Insertion at codon 69</li> </ul>	
d4T	<ul style="list-style-type: none"> <li>• V75A/M/S/T</li> <li>• T215A/C/D/E/G/H/I/L/N/S/V/Y/F [6]</li> <li>• At least 3 mutations among: M41L, D67N, K70R, L210W, K219Q/E [4, 7, 14, 15]</li> <li>• K65R [30, 31, 32]</li> <li>• Q151M</li> <li>• Insertion at codon 69</li> </ul>	
ABC	<ul style="list-style-type: none"> <li>• At least 3 mutations among: M41L, D67N, M184V/I, L210W, T215A/C/D/E/G/H/I/L/N/S/V/Y/F [8, 19, 29]</li> <li>• K65R [9, 11, 12]</li> <li>• L74V/I [24, 25, 26, 27, 28, 29]</li> <li>• Y115F</li> <li>• Q151M</li> <li>• Insertion at codon 69</li> </ul>	<ul style="list-style-type: none"> <li>• 2 mutations among: M41L, D67N, L210W, T215A/C/D/E/G/H/I/L/N/S/V/Y/F [8, 19, 29]</li> <li>• M184V/I [36]</li> </ul>
TDF/TAF	<ul style="list-style-type: none"> <li>• At least 4 mutations among: M41L, E44D, D67N, T69D/N/S, L74V/I, L210W, T215A/C/D/E/G/H/I/L/N/S/V/Y/F [13, 20, 33, 37, 38]</li> <li>• K65R/E/N [9, 10, 11, 12, 34, 35, 37, 38]</li> <li>• Insertion at codon 69</li> <li>• K70E [21, 22, 23]</li> </ul>	<ul style="list-style-type: none"> <li>• 3 mutations among: M41L, E44D, D67N, T69D/N/S, L74V/I, L210W, T215A/C/D/E/G/H/I/L/N/S/V/Y/F [13, 33, 37, 38]</li> </ul>

ZDV: zidovudine, 3TC: lamivudine, FTC: emtricitabine, ddl: didanosine, d4T: stavudine, ABC: abacavir, TDF: tenofovir, TAF : tenofovir alafenamide

**ANRS - AC 11: RESISTANCE GROUP  
GENOTYPE INTERPRETATION: NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS**

	Mutations associated with resistance	Mutations associated with « possible resistance »
EFV	<ul style="list-style-type: none"> <li>• L100I</li> <li>• K101E</li> <li>• K103H/N/S/T [1]</li> <li>• V106M [2]</li> <li>• E138K [12, 13]</li> <li>• Y181C/I</li> <li>• Y188C/L</li> <li>• G190A/C/E/Q/S/T/V</li> <li>• P225H</li> <li>• M230L</li> </ul>	
NVP	<ul style="list-style-type: none"> <li>• A98S (for HIV-1 subtype C only) [3]</li> <li>• L100I</li> <li>• K101E</li> <li>• K103H/N/S/T [1]</li> <li>• V106A/M [2]</li> <li>• Y181C/I</li> <li>• Y188C/H/L</li> <li>• G190A/C/E/Q/S/T/V</li> <li>• M230L</li> </ul>	<ul style="list-style-type: none"> <li>• E138K [13]</li> </ul>
ETR	<ul style="list-style-type: none"> <li>• At least 3 among: V90I, A98G, L100I, K101E/H/I/P/R, V106I, V179D/F/I/L/M/T, G190A/S, M230L [4, 7, 8, 9, 10, 11]</li> <li>• E138K [12, 13]</li> <li>• Y181C/I/V [5, 6]</li> <li>• H221Y [12,16]</li> </ul>	<ul style="list-style-type: none"> <li>• 2 mutations among: V90I, A98G, L100I, K101E/H/I/P/R, V106I, V179D/F/I/L/M/T, G190A/S, M230L [4, 7, 8, 9, 10, 11]</li> <li>• E138A/G/Q/R/S [5, 6, 7, 8]</li> </ul>
RPV	<ul style="list-style-type: none"> <li>• K101E/P [9, 13]</li> <li>• E138A/G/K/Q/R/S [12, 13, 14]</li> <li>• V179L [9]</li> <li>• Y181C/I/V [13]</li> <li>• Y188L [9]</li> <li>• H221Y [13]</li> <li>• M230I/L/V [9]</li> <li>• L100I + K103N/S [9, 15]</li> <li>• L100I + K103R + V179D [15]</li> </ul>	<ul style="list-style-type: none"> <li>• V179D [9,15, 17]</li> </ul>

EFV: efavirenz, NVP: nevirapine, ETR: etravirine, RPV : rilpivirine

**ANRS - AC 11: RESISTANCE GROUP  
GENOTYPE INTERPRETATION: PROTEASE INHIBITORS**

	Mutations associated with resistance	Mutations associated with « possible resistance »
IDV	<ul style="list-style-type: none"> <li>• M46I/L</li> <li>• V82A/F/M/S/T [11]</li> <li>• I84A/V [8]</li> <li>• L90M and at least 2 among: K20M/R, L24I, V32I, M36I, I54V/L/M/T, A71V/T, G73S/A, V77I</li> </ul>	<ul style="list-style-type: none"> <li>• L90M</li> </ul>
SQV/RTV 1000/100 mg BID	<ul style="list-style-type: none"> <li>• G48V</li> <li>• At least 3 mutations among: L10F/I/M/R/V, I15A/V, K20I/M/R/T, L24I, I62V, G73S/T, V82A/F/S/T, I84V, L90M [9]</li> </ul>	<ul style="list-style-type: none"> <li>• 2 mutations among: L10F/I/M/R/V, I15A/V, K20I/M/R/T, L24I, I62V, G73S/T, V82A/F/S/T, I84V, L90M [9]</li> </ul>
NFV	<ul style="list-style-type: none"> <li>• D30N</li> <li>• I84A/V [8]</li> <li>• N88S/D</li> <li>• L90M</li> </ul>	<ul style="list-style-type: none"> <li>• V82A/F/S/T and at least 2 among: L10I, M36I, M46I/L, I54V/L/M/T, A71V/T, V77I [1]</li> </ul>
LPV/r	<ul style="list-style-type: none"> <li>• At least 4 mutations among: L10F/I/R/V, K20M/R, L24I, L33F, M46I/L, I50V, F53L, I54M/L/T/V, L63P, A71I/L/V/T, V82A/F/S/T, I84V, L90M [3, 4, 5, 21]</li> <li>• I47A [15, 16]</li> <li>• L76V [18, 19]</li> </ul>	<ul style="list-style-type: none"> <li>• 3 mutations among: L10F/I/R/V, K20M/R, L24I, L33F, M46I/L, I50V, F53L, I54M/L/T/V, L63P, A71I/L/V/T, V82A/F/S/T, I84V, L90M [3, 4, 5, 21]</li> </ul>
FPV/RTV 700/100 mg BID	<ul style="list-style-type: none"> <li>• I50V</li> <li>• V32I and I47A/V [2, 13, 14]</li> <li>• At least 4 mutations among: L10F/I/V, L33F, M36I, I54A/L/M/S/T/V, I62V, V82A/C/F/G, I84V, L90M [2, 20]</li> </ul>	
ATV/RTV 300/100 mg QD	<ul style="list-style-type: none"> <li>• I50L [6]</li> <li>• N88S [28,29,30]</li> <li>• At least 3 mutations among: L10F/I/V, G16E, L33F/I/V, M46I/L, D60E, A71V/T, I84V, I85V, L90M [7, 12, 22, 31]</li> </ul>	<ul style="list-style-type: none"> <li>• 2 mutations among: L10F/I/V, G16E, L33F/I/V, M46I/L, D60E, A71V/T, I84V, I85V, L90M [7, 12, 22, 31]</li> </ul>
TPV/RTV 500/200 mg BID	<ul style="list-style-type: none"> <li>• At least a score of + 3*: 36I/L/V – 53L/W/Y + 58E + 69I/K/N/Q/R/Y + 89I/M/R/T/V [10, 23]</li> </ul>	<ul style="list-style-type: none"> <li>• A score of + 2*: 36I/L/V – 53L/W/Y + 58E + 69I/K/N/Q/R/Y + 89I/M/R/T/V [10, 23]</li> </ul>

September 2016- Version n°26

<b>DRV/RTV**</b> <b>600/100 mg BID</b>	<ul style="list-style-type: none"><li>• At least 4 mutations among: V11I, V32I, L33F, I47V, I50V, I54L/M, T74P, L76V, I84V, L89V [17, 24, 25, 26, 27]</li></ul>	<ul style="list-style-type: none"><li>• 3 mutations among: V11I, V32I, L33F, I47V, I50V, I54L/M, T74P, L76V, I84V, L89V [17, 24, 25, 26, 27]</li></ul>
<b>800/100 mg QD</b>	<ul style="list-style-type: none"><li>• 2 mutations among: V11I, V32I, L33F, I47V, I50V, I54L/M, T74P, L76V, I84V, L89V [17, 24, 25, 26, 27]</li></ul>	

IDV: indinavir, SQV: saquinavir, NFV: nelfinavir, RTV: ritonavir, FPV: fosamprenavir, LPV: lopinavir, ATV:atazanavir, TPV: tipranavir, DRV : darunavir

\* Insufficient data for HIV-1 subtype non-B

\*\* Please note that rules are different for DRV/RTV 600/100 mg BID and 800/100 mg QD

ANRS - AC 11: RESISTANCE GROUP

GENOTYPE INTERPRETATION: FUSION INHIBITOR

	Mutations associated with resistance
ENF	<ul style="list-style-type: none"><li>• G36A/D/E/S/V [1, 2, 3, 4, 5, 6, 7]</li><li>• V38A/E/K/M</li><li>• Q40H/K/P/T</li><li>• N42D/T</li><li>• N43D/H/K/S</li><li>• L44M</li><li>• L45Q/M</li></ul>
T20	

ENF (T20): enfuvirtide

**ANRS - AC 11 : RESISTANCE GROUP**  
**GENOTYPE INTERPRETATION: INTEGRASE STRAND TRANSFER INHIBITORS**

	<b>Mutations associated with resistance</b>	<b>Mutations associated with « possible resistance »</b>
<b>RAL</b>	<ul style="list-style-type: none"> <li>• T66K [10]</li> <li>• E92Q [1, 2]</li> <li>• G118R [10, 17]</li> <li>• F121Y [10,17]</li> <li>• G140A/S [7]</li> <li>• Y143A/C/G/H/R/S [1, 3, 4, 5, 8, 14]</li> <li>• Q148E/G/H/K/R [1, 2]</li> <li>• V151L [9]</li> <li>• N155H/S/T [1, 2, 9]</li> <li>• E157Q [2]</li> <li>• A49G + S230G/R + R263K [18]</li> </ul>	
<b>EVG</b>	<ul style="list-style-type: none"> <li>• T66I/A/K [6]</li> <li>• E92Q [6]</li> <li>• T97A [21,22]</li> <li>• G118R [17]</li> <li>• F121Y [9,17]</li> <li>• E138K</li> <li>• G140C/S</li> <li>• Y143A/C/G/H/R/S [14]</li> <li>• P145S [9]</li> <li>• S147G [21]</li> <li>• Q148H/R/K [6]</li> <li>• V151L [9]</li> <li>• N155H/S/T [6,9]</li> <li>• E157Q [11]</li> <li>• R263K [18]</li> </ul>	

<p>DTG* 50 mg BID</p>	<ul style="list-style-type: none"> <li>• G118R [12,13]</li> <li>• F121Y [17]</li> <li>• V151L [9]</li> <li>• S153Y [9]</li> <li>• R263K [16]</li> <li>• T66K + L74M [9]</li> <li>• E92Q + N155H [9, 23, 24]</li> <li>• Q148H/K/R + at least 2 mutations among: L74I or E138A/K/T or G140A/C/S [15]</li> <li>• Q148R + N155H [9]</li> </ul>	<ul style="list-style-type: none"> <li>• T66K [9]</li> <li>• S153F [9]</li> <li>• E157Q [19, 20]</li> <li>• Q148H/K/R + 1 mutation among: L74I or E138A/K/T or G140A/C/S [15]</li> </ul>
<p>50 mg QD</p>	<ul style="list-style-type: none"> <li>• G118R [12,13]</li> <li>• F121Y [17]</li> <li>• E138A/K/T</li> <li>• G140A/C/S</li> <li>• Q148H/K/R</li> <li>• V151L [9]</li> <li>• S153Y [9]</li> <li>• N155H [18]</li> <li>• R263K [16]</li> <li>• T66K + L74M [9]</li> </ul>	<ul style="list-style-type: none"> <li>• T66K [9]</li> <li>• S153F [9]</li> <li>• E157Q [19, 20]</li> </ul>
<p>CBG</p>	<ul style="list-style-type: none"> <li>• Q148K/R [25, 26, 27]</li> </ul>	

RAL: raltegravir, EVG: elvitegravir, DTG: dolutegravir, CBG: cabotegravir

\* Please note that rules are different for DTG 50 mg BID and 50 mg QD

## REFERENCES

### Nucleoside and Nucleotide Reverse Transcriptase Inhibitors

- 1/ Riva C et al. Transmitted virus with substitutions at position 215 and risk of virological failure in antiretroviral naive patients starting highly active antiretroviral therapy. XI International HIV drug resistance workshop : basic principles and clinical implications , 2-5 July 2002, Seville, Spain, abstract 124.
- 2/ Lanier ER et al. Effect of baseline 215D/C/S « revertant » mutations on virological response to lamivudine /zidovudine- containing regimens and emergence of 215Y upon virological failure. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain abstract 146.
- 3/ Garcia-Lerma et al. Increased ability for selection of zidovudine resistance in a distinct class of wild-type HIV-1 from drug-naive persons. PNAS 2001 ; 98 : 13907-13912.
- 4/ Chappey C et al. Evolution of amino acid 215 in HIV-1 reverse transcriptase in response to intermittent drug selection. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 32.
- 5/ Costagliola D et al. Presence of thymidine –associated mutations and response to d4T, abacavir and ddi in the control arm of the Narval ANRS 088 trial. 8<sup>th</sup> Conference on Retroviruses and opportunistic Infections. 4-8 february 2001, Chicago, Illinois, poster 450.
- 6/ Calvez V et al. Impact of stavudine phenotype and thymidine analogs mutations on viral response to stavudine plus lamivudine in ALTIS 2 ANRS trial. Antiviral Therapy 2002, 7(3):211-218.
- 7/ García-Lerma JG, MacInnes H, Bennett D, Reid P, Nidtha S, Weinstock H, Kaplan JE, Heneine W. A novel genetic pathway of human immunodeficiency virus type 1 resistance to stavudine mediated by the K65R mutation. J Virol. 2003 May;77(10):5685-93.
- 8/ Brun-Vézinet F et al. Clinically relevant interpretation of genotype for resistance to abacavir : a study from the Narval trial (ANRS 088). AIDS 2003; 17(12):1795-802.
- 9/ Stone C, Ait-Khaled M, Craig C, Griffin P, Tisdale M. Human immunodeficiency virus type 1 reverse transcriptase mutation selection during in vitro exposure to tenofovir alone or combined with abacavir or lamivudine. Antimicrob Agents Chemother. 2004 Apr;48(4):1413-5.
- 10/ Miller MD et al. Multivariate analyses of antiviral response to tenofovir DF therapy in antiretroviral–experienced patients. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain abstract 14.
- 11/ Margot NA, Lu B, Cheng A, Miller MD; Study 903 Team. Resistance development over 144 weeks in treatment-naive patients receiving tenofovir disoproxil fumarate or stavudine with lamivudine and efavirenz in Study 903. HIV Med. 2006 Oct;7(7):442-50.
- 12/ Parikh et al. K65R : a multi-nucleoside resistance mutation of a low but increasing frequency. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 136.

## September 2016- Version n°26

- 13/ Masquelier B et al. Genotypic and pharmacological determinants of the virological response to tenofovir in nucleoside reverse transcriptase inhibitor-experienced patients. *Antivir Ther.* 2004 ; 9(3):315-23.
- 14/ Izopet J. et al. Mutations conferring resistance to zidovudine diminish the antiviral effect of stavudine plus didanosine. *Journal of Medical Virology* 1999 ; 59 : 507-511.
- 15/ Pellegrin I. et al. Emergence of zidovudine and multidrug-resistance mutations in the HIV-1 reverse transcriptase gene in therapy-naive patients receiving stavudine plus didanosine combination therapy. STADI group. *AIDS* 1999 ; 13 : 1705-1709.
- 16/ Mulamba GB et al. Pre-steady state kinetic analysis of the incorporation of FTC 5'-monophosphate and 3TC 5'-monophosphate by mutants HIV-1 RTs K65R, K65R/Q151M and Q151M. 16<sup>th</sup> International Conference on Antiviral Research, 27 April-1 May 2003, Savannah, USA. Abstract 39.
- 17/ Marcelin AG et al. Clinically relevant genotype interpretation of resistance to didanosine. *Antimicrob Agents Chemother.* 2005 May;49(5):1739-44.
- 18/ Capdepon S et al. An additive/subtractive genotypic score as a determinant of the virological response to didanosine in HIV-1 infected patients. *J Clin Virol.* 2006 May;36(1): 36-42.
- 19/ Assoumou L, Brun-Vézinet F, Cozzi-Lepri A, Kuritzkes D, Phillips A, Zolopa A, Degruittola V, Miller V, Costagliola D; Standardization and Clinical Relevance of HIV Drug Resistance Testing Project of the Forum for Collaborative HIV Research. Initiatives for Developing and Comparing Genotype Interpretation Systems: External Validation of Existing Systems for Didanosine against Virological Response. *J Infect Dis.* 2008 Jul 3.
- 20/ Waters J et al. K65R, L74V and TAMs in HIV-1 RT associated with reduced response to tenofovir DF in antiretroviral-experienced patients. 13<sup>th</sup> Conference on Retroviruses and Opportunistic Infections. 5-8 february 2006, Denver, Colorado, poster 633.
- 21/ Kagan RM, Lee TS, Ross L, Lloyd RM Jr, Lewinski MA, Potts SJ. Molecular basis of antagonism between K70E and K65R tenofovir-associated mutations in HIV-1 reverse transcriptase. *Antiviral Res.* 2007 Sep;75(3):210-8.
- 22/ Kagan R et al. Adefovir HIV-1 RT mutation K70E in the age of tenofovir. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 93.
- 23/ Sluis-Cremer N, Sheen CW, Zelina S, Torres PS, Parikh UM, Mellors JW. Molecular mechanism by which the K70E mutation in human immunodeficiency virus type 1 reverse transcriptase confers resistance to nucleoside reverse transcriptase inhibitors. *Antimicrob Agents Chemother.* 2007 Jan;51(1):48-53.
- 24/ Miller V et al. HIV-1 reverse transcriptase (RT) genotype and susceptibility to RT inhibitors during abacavir monotherapy and combination therapy. *AIDS* 2000; 14:163–171.
- 25/ Moyle GJ, et al. Ziagen Once-Daily in Antiretroviral Combination Therapy (CNA30021) Study Team. Abacavir once or twice daily combined with once-daily lamivudine and efavirenz for the treatment of antiretroviral-naive HIV-infected adults: results of the Ziagen Once Daily in Antiretroviral Combination Study. *J Acquir Immune Defic Syndr.* 2005 Apr 1;38(4):417-25.

## September 2016- Version n°26

- 26/ Irlbeck D *et al.* Treatment-emergent mutations for previously naïve HIV-infected adults failing ZDV+3TC+EFV and ABC+3TC+EFV (CNA30024). 11<sup>th</sup> Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, USA. February 8–11, 2004. Abstract 661.
- 27/ Rodriguez-French A, *et al.* The NEAT study: a 48-week open-label study to compare the antiviral efficacy and safety of GW433908 versus nelfinavir in antiretroviral therapy-naïve HIV-1-infected patients. *J Acquir Immune Defic Syndr* 2004; **35**:22–32.
- 28/ Gathe JC Jr *et al.* SOLO: 48-week efficacy and safety comparison of once-daily fosamprenavir /ritonavir versus twice-daily nelfinavir in naïve HIV-1-infected patients. *AIDS*. 2004 Jul 23;18(11):1529-37
- 29/ Wirden *et al.* Risk factors for selection of the L74I reverse transcriptase mutation in human immunodeficiency virus type 1-infected patients. *Antimicrob Agents Chemother*. 2006 Jul;50(7):2553-6.
- 30/García-Lerma JG *et al.* A novel genetic pathway of human immunodeficiency virus type 1 resistance to stavudine mediated by the K65R mutation. *J Virol*. 2003 May;77(10):5685-93
- 31/Sungkanuparph S *et al.* Prevalence and risk factors for developing K65R mutations among HIV-1 infected patients who fail an initial regimen of fixed-dose combination of stavudine, lamivudine, and nevirapine. *J Clin Virol*. 2008 ;41:310-3.
- 32/Coutsinos D *et al.* The K65R mutation subtype B and C HIV-1: rates of development and the implications of template-specific dislocation mutagenesis. *Antiviral Therapy* 2010; 15 suppl 2:A13 (abs 5)
- 33/ Wirden M *et al.*, Antiretroviral combinations implicated in emergence of the L74I and L74V resistance mutations in HIV-1-infected patients. *AIDS*. 2009 Jan 2;23(1):95-9.
- 34/ Fourati S *et al.* Identification of a rare mutation at reverse transcriptase Lys65 (K65E) in HIV-1-infected patients failing on nucleos(t)ide reverse transcriptase inhibitors. *J Antimicrob Chemother*. 2013 Jun 19.
- 35/ Ross LL *et al.* A rare HIV reverse transcriptase mutation, K65N, confers reduced susceptibility to tenofovir, lamivudine and didanosine. *AIDS* 2006 Mar 21;20(5):787-9.
- 36/ Tisdale M *et al.* Combination of mutations in human immunodeficiency virus type 1 reverse transcriptase required for resistance to the carbocyclic nucleoside 1592U89. *Antimicrob Agents Chemother*. 1997 May;41(5):1094-8.
- 37/Callebaut C *et al.* In Vitro Virology Profile of Tenofovir Alafenamide, a Novel Oral Prodrug of Tenofovir with Improved Antiviral Activity Compared to That of Tenofovir Disoproxil Fumarate. *Antimicrob Agents Chemother*. 2015 Oct;59(10):5909-16.
- 38/Margot NA *et al.* Rare emergence of drug resistance in HIV-1 treatment-naïve patients after 48 weeks of treatment with elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide. *HIV Clin Trials*. 2016 Mar;17(2):78-87.

**Non nucleoside transcriptase inhibitors**

- 1/ Harrigan PR, Mo T, Wynhoven B, Hirsch J, Brumme Z, McKenna P, Pattery T, Vingerhoets J, Bachelier LT. Rare mutations at codon 103 of HIV-1 reverse transcriptase can confer resistance to non-nucleoside reverse transcriptase inhibitors. *AIDS*. 2005 Mar 24;19(6):549-54.
- 2/ Brenner B et al. A V106M mutation in HIV-1 clade C viruses exposed to efavirenz confers cross-resistance to non-nucleoside reverse transcriptase inhibitors. *AIDS*. 2003 Jan 3;17(1):F1-5.
- 3/ Deshpande A et al. Resistance mutations in subtype C HIV type 1 isolates from Indian patients of Mumbai receiving NRTIs plus NNRTIs and experiencing a treatment failure: resistance to AR. *AIDS Res Hum Retroviruses* 2007; 23 : 335-40.
- 4/ Vingerhoets J et al. Impact of baseline NNRTI mutations on the virological response to TMC125 in the Phase III clinical trials DUET-1 and DUET-2. XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 32.
- 5/ Marcelin AG et al. Factors associated with virological response to etravirine in nonnucleoside reverse transcriptase inhibitor-experienced HIV-1-infected patients. *Antimicrob Agents Chemother*. 2010 Jan;54(1):72-7.
- 6/ Vingerhoets J et al. Resistance profile of etravirine: combined analysis of baseline genotypic and phenotypic data from the randomized, controlled Phase III clinical studies. *AIDS*. 2010 Feb 20;24(4):503-14.
- 7/ Vingerhoets J et al. Impact of transmitted drug resistance (TDR) on virological response to initial combination antiretroviral therapy (cART). *Antiviral Therapy* 2010; 15 Suppl 2: A125 (abstract 99).
- 8/ El A et al. In vitro selection of novel etravirine-associated resistance mutations in B and non-B HIV-1 subtypes. *Antiviral Therapy* 2010; 15 Suppl 2: A134 (abstract 108).
- 9/ Azijn H et al. TMC278, a next-generation nonnucleoside reverse transcriptase inhibitor (NNRTI), active against wild-type and NNRTI-resistant HIV-1. *Antimicrob Agents Chemother*. 2010 Feb;54(2):718-27.
- 10/ Marcelin AG et al. Mutations selected in patients displaying treatment failure under an etravirine-containing regimen. *Antiviral Therapy* 2011; 10.3851/IMP1886.
- 11/ Maïga AI et al. Resistance-associated mutations to etravirine (TMC-125) in antiretroviral-naïve patients infected with non-B HIV-1 subtypes. *Antimicrob Agents Chemother*. 2010 Feb;54(2):728-33.
- 12/ Asahchop EL et al. Characterization of the E138K resistance mutation in HIV-1 reverse transcriptase conferring susceptibility to etravirine in B and non-B HIV-1 subtypes. *Antimicrob Agents Chemother*. 2011 Feb;55(2):600-7.

## September 2016- Version n°26

13/ Eron J et al. Characterization of the resistance profile of TMC278: 48-week analysis of the Phase III studies ECHO and THRIVE. 50th ICAAC 2010 (abstract H-1810).

14/ Haddad M et al. Impact of HIV-1 reverse transcriptase E138 mutations selected on rilpivirine drug susceptibility. *Antiviral Therapy* 2011; 16 Suppl 1: A18 (abstract 10).

15/ Haddad M et al. Combinations of HIV-1 reverse transcriptase mutations L100I+K103N/S and L100I+K103R+V179D reduce susceptibility to rilpivirine. 53rd ICAAC 2013 (abstract H-677)

16/ Melikian et al. Non-nucleoside reverse transcriptase inhibitor (NNRTI) cross-resistance: implications for preclinical evaluation of novel NNRTIs and clinical genotypic resistance testing. *J Antimicrob Chemother.* 2014 Jan;69(1):12-20.

17/ Porter et al. Characterization of HIV-1 drug resistance development through week 48 in antiretroviral naive subjects on rilpivirine/emtricitabine/tenofovir DF or efavirenz/emtricitabine/tenofovir DF in the STaR study (GS-US-264-0110). *J Acquir Immune Defic Syndr.* 2014 Mar 1;65(3):318-26.

### Protease inhibitors

1/ Phenotype/genotype correlation from Stanford database.

2/ Masquelier B, Assoumou KL, Descamps D, Bocket L, Cottalorda J, Ruffault A, Marcelin AG, Morand-Joubert L, Tamalet C, Charpentier C, Peytavin G, Antoun Z, Brun-Vézinet F, Costagliola D; ANRS Resistance Study Group. Clinically validated mutation scores for HIV-1 resistance to fosamprenavir/ritonavir. *J Antimicrob Chemother.* 2008 Jun;61(6):1362-8.

3/ Marcelin AG, Cohen-Codar I, King MS, Colson P, Guillevic E, Descamps D, Lamotte C, Schneider V, Ritter J, Segondy M, Peigue-Lafeuille H, Morand-Joubert L, Schmuck A, Ruffault A, Palmer P, Chaix ML, Mackiewicz V, Brodard V, Izopet J, Cottalorda J, Kohli E, Chauvin JP, Kempf DJ, Peytavin G, Calvez V. Virological and pharmacological parameters predicting the response to lopinavir-ritonavir in heavily protease inhibitor-experienced patients. *Antimicrob Agents Chemother.* 2005 May;49(5):1720-6.

4/ Maillard A, Chaplain JM, Tribut O, Bentué-Ferrer D, Tattevin P, Arvieux C, Michelet C, Ruffault A. The use of drug resistance algorithms and genotypic inhibitory quotient in prediction of lopinavir-ritonavir treatment response in human immunodeficiency virus type 1 protease inhibitor-experienced patients. *J Clin Virol.* 2007 Feb;38(2):131-8.

5/ Masquelier B et al. Human Immunodeficiency virus type 1 genotypic and pharmacokinetic determinants of the virological response to lopinavir-ritonavir-containing therapy in protease inhibitor-experienced patients. *Antimicrob Agents and Chemother* 2002 ; 46 : 2926-2932.

## September 2016- Version n°26

- 6/ Colonna RJ et al. Identification of I50L as the signature atazanavir (ATV)-resistance mutation in treatment naïve HIV-1 infected patients receiving ATV-containing regimens. *Journal of Infectious Diseases* 2004; 189: 1802-10.
- 7/ Vora S et al. Clinical validation of atazanavir/ritonavir genotypic resistance score in PI-experienced patients. *AIDS* 2006 Jan 2;20(1):35-40.
- 8/ Mo H, Parkin N, Stewart KD, Lu L, Dekhtyar T, Kempf DJ, Molla A. Identification and structural characterization of I84C and I84A mutations that are associated with high-level resistance to human immunodeficiency virus protease inhibitors and impair viral replication. *Antimicrob Agents Chemother.* 2007 Feb;51(2):732-5.
- 9/ Marcelin AG, Flandre P, de Mendoza C, Roquebert B, Peytavin G, Valer L, Wirden M, Abbas S, Katlama C, Soriano V, Calvez V. Clinical validation of saquinavir/ritonavir genotypic resistance score in protease-inhibitor-experienced patients. *Antivir Ther.* 2007;12(2):247-52.
- 10/ Marcelin AG et al. Tipranavir-ritonavir genotypic resistance score in protease inhibitor-experienced patients. *Antimicrob Agents Chemother.* 2008 Sept;52(9).
- 11/ Camacho R et al. Different substitutions under selective pressure at protease codon 82 in HIV-1 subtype G compared to subtype B infected individuals including a novel I82M resistance mutation. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 138.
- 12/ Marcelin AG et al. External validation of atazanavir/ritonavir genotypic score in HIV-1 protease inhibitor experienced patients. *JAIDS* 2006; 42 (1): 127-8.
- 13/ Brutus A et al. The Neat Study: Virologic response in subjects receiving GW433908 (908) BID by viral load/CD4 cell Counts at study entry. XV International AIDS Conference, July 11-16 2004, Bangkok ,Thailand, abstract TuPeB 4500.
- 14/ Pellegrin I, Breilh D, Coureau G, Boucher S, Neau D, Merel P, Lacoste D, Fleury H, Saux MC, Pellegrin JL, Lazaro E, Dabis F, Thiébaud R; ANRS Co3 Aquitaine Cohort. Interpretation of genotype and pharmacokinetics for resistance to fosamprenavir-ritonavir-based regimens in antiretroviral-experienced patients. *Antimicrob Agents Chemother.* 2007 Apr;51(4):1473-80.
- 15/ Friend J et al. Isolated lopinavir resistance after virological rebound of a ritonavir/lopinavir-based regimen. *AIDS.* 2004 Sep 24;18(14):1965-6.
- 16/ de Mendoza C et al. Prevalence of the HIV-1 protease mutation I47A in clinical practice and association with lopinavir resistance. *AIDS* 2006 Apr 24; 20(7): 1071-4.
- 17/ de Meyer S, Vangeneugden T, van Baelen B, de Paepe E, van Marck H, Picchio G, Lefebvre E, de Béthune MP. Resistance profile of darunavir: combined 24-week results from the POWER trials. *AIDS Res Hum Retroviruses.* 2008 Mar;24(3):379-88.
- 18/ Nijhuis N et al. A novel genetic pathway involving L76V and M46I leading to lopinavir/r resistance. XVI International HIV drug resistance workshop : basic principles and clinical implications, 12-16 June 2007, Barbados, West Indies, abstract 127.
- 19/ Delaugerre C et al. Protease inhibitor resistance analysis in the MONARK trial comparing first-line lopinavir-ritonavir monotherapy to lopinavir-ritonavir plus zidovudine and lamivudine triple therapy. *Antimicrob Agents Chemother.* 2009 Jul;53(7):2934-9.

## September 2016- Version n°26

- 20/ Marcelin AG et al. Genotypic resistance analysis of the virological response to fosamprenavir-ritonavir in protease inhibitor-experienced patients in CONTEXT and TRIAD clinical trials. *Antimicrob Agents Chemother.* 2008 Dec;52(12):4251-7.
- 21/ Hill A et al. Identification of new genotypic cut-off levels to predict the efficacy of lopinavir/ritonavir and darunavir/ritonavir in the TITAN trial. *HIV Med.* 2009 Jul 6.
- 22/ Di Giambenedetto S et al. A rigorous statistical learning method for the estimation and validation of weighted drug susceptibility scores applied to in vivo virological outcome prediction in atazanavir/ritonavir-containing HAART. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 95.
- 23/ Bethell R et al. No effect of subtype on susceptibility and virological response to TPV/r for treatment experienced patients. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 111.
- 24/ Descamps D et al. Mutations associated with virological response to darunavir/ritonavir in HIV-1-infected protease inhibitor-experienced patients. *J Antimicrob Chemother.* 2009 Mar;63(3):585-92.
- 25/ De Meyer S. et al. Influence of baseline protease inhibitor resistance on the efficacy of darunavir/ritonavir or lopinavir/ritonavir in the TITAN trial. *J Acquir Immune Defic Syndr.* 2008 Dec 15;49(5):563-4.
- 26/ De Meyer S. et al. Phenotypic and genotypic determinants of resistance to darunavir: analysis of data from treatment-experienced patients in POWER 1, 2, 3 and DUET-1 and 2. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 31.
- 27/ De Meyer S. et al. Confirmation of the negative impact of protease mutations I47V, I54M, T74P and I84V and the positive impact of protease mutation V82A on virological response to darunavir/ritonavir. XVII International HIV drug resistance workshop : basic principles and clinical implications, 9-13 June 2009, Fort Myers, Florida, abstract 126.
- 28/ Gong YF. Et al. In vitro resistance profile of the human immunodeficiency virus type 1 protease inhibitor BMS-232632. *Antimicrob Agents Chemother.* 2000 Sep;44(9):2319-26.
- 29/ Malan DR; et al. Efficacy and safety of atazanavir, with or without ritonavir, as part of once-daily highly active antiretroviral therapy regimens in antiretroviral-naïve patients. *J Acquir Immune Defic Syndr.* 2008 Feb 1;47(2):161-7.
- 30/ Malan DR et al. 96-week efficacy and safety of atazanavir, with and without ritonavir, in a HAART regimen in treatment-naïve patients. *J Int Assoc Physicians AIDS Care (Chic).* 2010 Jan-Feb;9(1):34-42.
- 31/ Morand-Joubert L et al. 15<sup>th</sup> Emerging resistance mutations in protease inhibitor naïve patients failing atazanavir based regimen (ANRS multicentre observational study). European AIDs Conference. October 21-24, Barcelona, Spain (Abstract PE9/9)

**Fusion inhibitor**

1/ Sista P et al. Subgroup analysis of baseline susceptibility and early virological response to enfuvirtide in the combined TORO studies. XII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2003, Los Cabos, Mexico, abstract 55.

2/ Mink M et al. Impact of HIV-1 gp41 amino acid substitutions (position 36-45) on susceptibility to T20 (enfuvirtide) in vitro ; analysis of primary virus isolates recovered from patients during chronic enfuvirtide treatment and site-directed mutants in NL4-3. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain, abstract 22.

3/ Greenberg ML et al. Enfuvirtide (T-20) and T-1249 resistance : observations from phase II clinical trials of enfuvirtide in combination with oral antiretrovirals and a phase I/II dose-ranging monotherapy trial of T-1249. XI International HIV drug resistance workshop : basic principles and clinical implications, 2-5 July 2002, Seville, Spain, abstract 128.

4/ Greenberg ML et al. Baseline and on-treatment susceptibility to enfuvirtide seen in TORO 1 and TORO 2 to 24 weeks. 10<sup>th</sup> Conference on Retroviruses and Opportunistic Infections, February 10-14, 2003, Boston, USA, abstract 141.

5/ Chakraborty B et al. Replicative fitness of HIV-1 strains with reduced susceptibility to protease-, reverse transcriptase- and entry (enfuvirtide)-inhibitors. XIII International HIV drug resistance workshop : basic principles and clinical implications, 8-12 June 2004, Tenerife, Spain, abstract 61.

6/ Walmsley SL et al. Development of a new genotypic resistance assay involving the entire gp41 sequence for evaluating resistance to enfuvirtide. XIII International HIV drug resistance workshop : basic principles and clinical implications, 8-12 June 2004, Tenerife, Spain, abstract 134.

7/ Melby T et al. Evolution of enfuvirtide resistance in longitudinal samples obtained after continued enfuvirtide dosing post-virological failure. XIV International HIV Drug Resistance Workshop: basic principles and clinical implications, 7-11 June 2005, Québec City, Québec, Canada, abstract 67.

**Integrase strand transfer inhibitors**

1/ Cooper DA et al. Subgroup and resistance analyses of raltegravir for resistant HIV-1 infection. N Engl J Med. 2008 Jul 24;359(4):355-65.

2/ Malet I et al. Mutations associated with failure of raltegravir treatment affect integrase sensitivity to the inhibitor in vitro. Antimicrob Agents Chemother. 2008 Apr;52(4):1351-8.

3/ Hatano H et al. Evolution of integrase resistance during failure of integrase inhibitor-based antiretroviral therapy. J Acquir Immune Defic Syndr. 2010 Aug 1;54(4):389-93.

## September 2016- Version n°26

- 4/ Da Silva D et al. HIV-1 resistance patterns to integrase inhibitors in antiretroviral-experienced patients with virological failure on raltegravir-containing regimens. *J Antimicrob Chemother.* 2010 Jun;65(6):1262-9
- 5/ Ceccherini-Silberstein F et al. Virological response and resistance in multi-experienced patients treated by raltegravir. XVII International HIV drug resistance workshop : basic principles and clinical implications, 10-14 June 2008, Sitges, Spain, abstract 18.
- 6/ Waters J et al. Evolution of resistance to the HIV integrase inhibitor (INI) elvitegravir can involve genotypic switching among primary INI resistance patterns. XVII International HIV drug resistance workshop : basic principles and clinical implications, 9-13 June 2009, Fort Myers, Florida, abstract 116.
- 7/ Geretti AM et al. Prevalence and patterns of raltegravir resistance in treated patients in Europe. *Antiviral Therapy* 2010; 15 Suppl 2: A62 (abstract 51)
- 8/ Huang W et al. Identification of alternative amino acid substitutions at HIV-1 integrase codon 143 that confer reduced susceptibility to raltegravir. 18th Conference on Retroviruses and Opportunistic Infections, February 27-March 2, 2011, Boston, USA, abstract 607.
- 9/ Kobayashi M et al. In Vitro antiretroviral properties of S/GSK1349572, a next-generation HIV integrase inhibitor. *Antimicrob Agents Chemother.* 2011 Feb;55(2):813-21.
- 10/ Malet I et al. The HIV-1 integrase G118R mutation confers raltegravir resistance to the CRF02\_AG HIV-1 subtype. *J Antimicrob Chemother.* 2011 Dec;66(12):2827-30.
- 11/ White K et al. Integrated Analysis of Emergent Drug Resistance from the HIV-1 Phase 3 QUAD Studies through Week 48. International Workshop on HIV & Hepatitis Virus Drug Resistance and Curative Strategies June 5-9, 2012 Sitges, Spain, Abstract 4.
- 12/ Bar-Magen et al. Identification of novel mutations responsible for resistance to MK-2048, a second-generation HIV-1 integrase inhibitor. *J Virol.* 2010 Sep;84(18):9210-6.
- 13/ Hare S et al. Structural and functional analyses of the second-generation integrase strand transfer inhibitor dolutegravir (S/GSK1349572). *Mol Pharmacol.* 2011 Oct;80(4):565-72.
- 14/ Huang W et al. Contribution of raltegravir selected secondary mutations to reduction in elvitegravir susceptibility of patient-derived HIV integrase containing Y143 mutations. International Workshop on HIV & Hepatitis Virus Drug Resistance and Curative Strategies June 4-8, 2013 Toronto, Canada, Abstract 89.
- 15/ Vavro CL et al. Integrase genotypic and phenotypic predictors of antiviral response to dolutegravir (DTG) in subjects with resistance to integrase inhibitors (INIs). International Workshop on HIV & Hepatitis Virus Drug Resistance and Curative Strategies June 4-8, 2013 Toronto, Canada, Abstract 29.
- 16/ Underwood MR et al. Analysis and characterization of treatment-emergent resistance in ART-experienced, integrase inhibitor-naive subjects with dolutegravir (DTG) versus raltegravir (RAL) in SAILING (ING111762). International Workshop on HIV & Hepatitis Virus Drug Resistance and Curative Strategies June 4-8, 2013 Toronto, Canada, Abstract 21.

## September 2016- Version n°26

- 17/ Malet I et al. New raltegravir resistance pathways induce broad cross-resistance to all currently used integrase inhibitors. *J Antimicrob Chemother.* 2014 Aug;69(8):2118-22.
- 18/ Underwood MR et al. Resistance Post Week 48 in ART-Experienced, Integrase Inhibitor-Naïve Subjects with Dolutegravir (DTG) vs. Raltegravir (RAL) in SAILING (ING111762). 13th European HIV & Hepatitis workshop, june 3-5 2015, Barcelona, Spain (Abstract n°6)
- 19/ Walmsley S, et al. Dolutegravir Plus Abacavir/Lamivudine for the Treatment of HIV-1 Infection in Antiretroviral Therapy-Naive Patients: Week 96 and Week 144 Results from the SINGLE Randomized Clinical Trial. *J Acquir Immune Defic Syndr.* 2015 Aug 7.
- 20/ Danion et al. Non-virological response to a dolutegravir-containing regimen in a patient harbouring a E157Q-mutated virus in the integrase region. *J Antimicrob Chemother.* 2015 Jun;70(6):1921-3.
- 21/ Molina et al. Efficacy and safety of once daily elvitegravir versus twice daily raltegravir in treatment-experienced patients with HIV-1 receiving a ritonavir-boosted protease inhibitor: randomised, double-blind, phase 3, non-inferiority study. *Lancet Infect Dis.* 2012 Jan;12(1):27-35.
- 22/ Sax et al. Co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus co-formulated efavirenz, emtricitabine, and tenofovir for initial treatment of HIV-1 infection: a randomised, double-blind, phase 3 trial, analysis of results after 48 weeks. *Lancet.* 2012 Jun 30;379(9835):2439-48.
- 23/ Underwood et al. The activity of the integrase inhibitor dolutegravir against HIV-1 variants isolated from raltegravir-treated adults. *J Acquir Immune Defic Syndr.* 2012 Nov 1;61(3):297-301.
- 24/ Frantzell et al. Dolutegravir resistance requires multiple primary mutation in HIV-1 integrase. CROI 2015. Abstract Number: 121.
- 25/ Yoshinoga et al. Advanced mechanistic studies of GSK1265744, a new HIV integrase inhibitor (INI) dosed by oral administration or long acting parenteral injection. *Antiviral Therapy* 2013 ; 18 Suppl 1: A30. Abstract 22)
- 26/ Margolis et al. 744 and Rilpivirine As Two Drug Oral Maintenance Therapy: LAI116482 (LATTE) Week 48 Results. CROI 2014. Abstract Number: 91LB
- 27/ Dudas et al. Characterization of NNRTI and INI Resistance Mutations Observed in a Study Subject on Oral Two-Drug Maintenance Therapy with 10 mg Cabotegravir + 25 mg Rilpivirine. IHDRW 2015 Seattle, WA, USA. Abstract 13.